


Aerosol and surface stability of HCoV-19 (SARS-CoV-2) compared to SARS-CoV-1

Summary:

1. Virus transmission via respiratory secretions in the form of droplets (>5 microns) or aerosols (<5 microns) appears to be likely.
2. Researchers found that viable virus could be detected in aerosols up to 3 hours post aerosolization, up to 4 hours on copper, up to 24 hours on cardboard and up to 2-3 days on plastic and stainless steel.
3. Results indicate that aerosol and fomite transmission of HCoV-19 is plausible, as the virus can remain viable in aerosols for multiple hours and on surfaces up to days.
4. There have been early indications that individuals infected with HCoV-19 may shed and transmit the virus while pre-symptomatic or asymptomatic.
5. The epidemiology of SARS-CoV-1 was dominated by nosocomial transmission and SARS-CoV was detected on variety of surfaces and objects in healthcare settings. HCoV-19 transmission is also occurring in hospital settings, with over 3000 reported cases of hospital-acquired infections. These cases highlight the vulnerability of healthcare settings for introduction and spread of HCoV-19. However, in contrast to SARS-CoV-1, most secondary transmission has been reported outside healthcare settings and widespread transmission in the community is being seen in several settings, such as households, workplace and group gatherings.



Researchers:

1. Laboratory of Virology, Division of Intramural Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Hamilton, MT, USA
2. Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA
3. Department of Ecology and Evolutionary Biology, University of California, Los Angeles, Los Angeles, CA, USA
4. Division of Viral Diseases, National Center for Immunization and Respiratory Diseases, Centers for Disease Control and Prevention, Atlanta, GA, USA.
5. Fogarty International Center, National Institutes of Health, Bethesda, MD, USA

Research supported by U.S. Department of Defense, Defense Advanced Research Projects Agency, U.S. National Science Foundation, Strategic Environmental Research and Development Program

[Link to Peer Reviewed Journal Study – Published March 17, 2020 in The New England Journal of Medicine](#)